


Proportion of Cancers Associated with Excess Body Weight Varies Considerably by State

Cancers related to body weight twice as predominant in women than in men

C Both sexes combined



A new study looking at the share of cancers related to obesity finds an at least 1.5-fold difference between states with the highest and lowest proportions. The proportion of cancer cases that could be attributable to excess body weight ranged from a high of 8.3% in the District of Columbia to a low of 5.9% in Hawaii, reflecting variations in obesity rates in the states. The study appears in [JAMA Oncology](#).

Excess body weight is an established cause of cancer, currently known to be linked to 13 cancers. While differences in excess body weight among states in the United

Rank and Proportion of Incident Cancer Cases in the 50 US States and District of Columbia Attributable to Excess Body Weight in Adults 30 Years or Older, 2011-2015

The figures show estimates based on the prevalence of excess body weight from 2001 to 2004. States are categorized using the Jenks natural breaks classification method.

States are well-known, there is little information on the burden of obesity-related cancers by state, information that would be useful for setting priorities for cancer control initiatives.

To learn more, American Cancer Society investigators led by Farhad Islami, M.D., Ph.D., Scientific Director, Surveillance Research, calculated the population attributable fraction (PAF) of incident cancer cases attributable to excess body weight among adults aged ≥ 30 years in 2011-2015 in all 50 states and the District of Columbia.

They found that in both men and women, there was at least a 1.5-fold difference between states with the highest and lowest proportions of cancers attributable to excess body weight. Among men, the population attributable fraction (PAF) ranged from 3.9% in Montana to 6.0% in Texas. Meanwhile, the PAF for women was approximately twice as high as for men, ranging from 7.1% in Hawaii to 11.4% in the District of Columbia. States with largest PAFs were mostly located in the South and Midwest, plus Alaska and the District of Columbia.

The proportion was far greater for some individual cancer types. For endometrial cancer, for example, the PAF ranged from 36.5% in Hawaii to 54.9% in Mississippi and was 50.0% or more in 19 states.

“The proportion of cancers attributable to [excess body weight] varies among states, but [excess body weight] accounts for at least 1 in 17 of all incident cancers in each state,” write the authors. “Broad implementation of known community- and individual-level interventions is needed to reduce access to and marketing of unhealthy foods (eg, through a tax on sugary drinks) and to promote and increase access to healthy foods and physical activity, as well as preventive care.”

“ *Broad implementation of known community- and individual-level interventions is needed to reduce access to and marketing of unhealthy foods... and to promote and increase access to healthy foods and physical activity* ”

Article: Proportion of cancer cases attributable to excess body weight by state in the United States, 2011-2015 JAMA Oncol 2018 DOI: 10.1001/jamaoncol.2018.5639