

American Cancer Society Updates Lung Cancer Screening Guideline: Nearly Five Million U.S. Adults who Smoke and Formerly Smoked Now Recommended for Testing

For the start of Lung Cancer Awareness Month, the American Cancer Society aims to reduce lung cancer deaths with updated screening recommendations

ATLANTA, November 1, 2023 — Today, the [American Cancer Society](#) (ACS) released an update of its lung cancer screening guideline to help reduce the number of people dying from the disease due to smoking history. The new guideline recommends yearly screening for lung cancer for people aged 50 to 80 years old who smoke or formerly smoked and have a 20-year or greater pack-year history. The recommended annual screening test for lung cancer is a low-dose computed tomography scan (also called a low-dose CT scan, or LDCT). The guideline, last updated in 2013, is published in the ACS flagship journal, [CA: A Cancer Journal for Clinicians](#).

“This updated guideline continues a trend of expanding eligibility for lung cancer screening in a way that will result in many more deaths prevented by expanding the eligibility criteria for screening to detect lung cancer early,” said [Dr. Robert Smith](#), senior vice president, early cancer detection science at the American Cancer Society and lead author of the lung cancer screening guideline report. “Recent studies have shown extending the screening age for persons who smoke and formerly smoked, eliminating the ‘years since quitting’ requirement and lowering the pack per year recommendation could make a real difference in saving lives.”

About Lung Cancer

Lung cancer is the overall leading cause of cancer death in the United States and is the second most frequently diagnosed malignancy in both men and women. In 2023, ACS researchers estimate 238,340 new cases of lung cancer (117,550 in men and 120,790 in women) will be diagnosed with about 127,070 deaths from the disease (67,160 in men and 59,910 in women). Anyone at any age can get lung cancer. However, lung cancer mainly occurs in older people, as most people diagnosed with the disease are aged 65 or older.

The updated guideline report is part of the ongoing guideline development process by ACS scientists. The ACS monitors medical and scientific literature for new evidence that may support a change in current guidelines or the development of a new guideline and information about screening that should be conveyed to clinicians and target populations.

How does the new guideline differ from the previously published guideline?

Item	Previous Recommendations	New Recommendations
Age for eligibility	55-74 years	50-80 years
Pack-year (PY) history	30+ PY	20+ PY
Years since quitting (YSQ)	≤ 15 YSQ	No Longer Required

In addition, report authors stress it’s important for people who are going to be screened to consult with their doctor about the potential benefits, limits, and harms of yearly screening with LDCT scans. People who still smoke should receive counseling to help them quit.

Who should NOT get screened for lung cancer:

Adults with health conditions that would greatly limit life expectancy or affect their ability or willingness to get lung cancer treatment if diagnosed should not be screened for the disease.

The American Cancer Society's advocacy affiliate, the American Cancer Society Cancer Action Network (ACS CAN) works at all levels of government to advocate for policies to expand access to lung cancer screenings and tobacco control.

"This updated guideline is critical to identify all individuals who can benefit from the early detection of lung cancer," said [Lisa Lacasse](#), president of ACS CAN. "ACS CAN will continue to work to improve access to care and lung cancer screening at no cost for any person at risk of the disease to decrease lung cancer deaths and help end cancer as we know it, for everyone."

"The good news is our research shows the number of new lung cancer cases diagnosed each year continues to decrease, partly because more people are quitting smoking (or not starting)," added Smith. "The number of deaths from lung cancer continues to drop as well, due to fewer people smoking and advances in early detection and treatment, but we still have to do better. This updated guideline is a step in the right direction."

Today's published guideline report also includes a [patient page](#), an [editorial from CA editors](#), and a [systematic review](#) of the evidence supporting dropping the years-since-quit criterion related to the new guideline. Modeling papers and epidemiologic research that support the new ACS guideline, as well as an updated analysis of the benefit-to-radiation-risk ratio from screening, are also published today in the ACS journal [Cancer](#).

Other ACS authors include: [Dr. Karli Kondo](#), [Tyler Kratzer](#), [Dr. Deana Baptiste](#), and [Dr. William Dahut](#).



The complete [ACS updated lung cancer screening guideline](#), and additional information about lung cancer prevention, diagnosis, and treatment is available on [cancer.org](#), including the [ACS Empowered to Quit program](#).

#

About the American Cancer Society

The American Cancer Society is a leading cancer-fighting organization with a vision to end cancer as we know it, for everyone. For more than 100 years, we have been improving the lives of people with cancer and their families as the only organization combating cancer through advocacy, research, and patient support. We are committed to ensuring everyone has an opportunity to prevent, detect, treat, and survive cancer. To learn more, visit [cancer.org](#) or call our 24/7 helpline at 1-800-227-2345. Connect with us on [Facebook](#), [Twitter](#), and [Instagram](#).

For further information: American Cancer Society, Anne.Doerr@cancer.org

Additional assets available online:  [Video \(3\)](#)  [Photos \(3\)](#)